

Pattern of self-reported oral disadvantage due to disease.

Baseline	6-month	12-month	18-month	24-month	Percent of Sample with Each Pattern (n=755)
No	No	No	No	No	65
No	No	No	No	Yes	3
No	No	No	Yes	No	1
No	No	No	Yes	Yes	0
No	No	Yes	No	No	1
No	No	Yes	No	Yes	0
No	No	Yes	Yes	No	1
No	No	Yes	Yes	Yes	0
No	Yes	No	No	No	2
No	Yes	No	No	Yes	0
No	Yes	No	Yes	No	0
No	Yes	No	Yes	Yes	0
No	Yes	Yes	No	No	0
No	Yes	Yes	No	Yes	0
No	Yes	Yes	Yes	No	0
No	Yes	Yes	Yes	Yes	1
Yes	No	No	No	No	5
Yes	No	No	No	Yes	3
Yes	No	No	Yes	No	0
Yes	No	No	Yes	Yes	1
Yes	No	Yes	No	No	1
Yes	No	Yes	No	Yes	1
Yes	No	Yes	Yes	No	1
Yes	No	Yes	Yes	Yes	1
Yes	Yes	No	No	No	2
Yes	Yes	No	No	Yes	1
Yes	Yes	No	Yes	No	0
Yes	Yes	No	Yes	Yes	1
Yes	Yes	Yes	No	No	1
Yes	Yes	Yes	No	Yes	1
Yes	Yes	Yes	Yes	No	1
Yes	Yes	Yes	Yes	Yes	5

Notes:

To eliminate permutations that include missing data, this aspect of the analysis is based on the 755 subjects who provided information on oral disadvantage at baseline and at every follow-up during the 24-month observation period.

Pattern of self-reported oral disadvantage due to pain.

Baseline	6-month	12-month	18-month	24-month	Percent of Sample with Each Pattern (n=752)
No	No	No	No	No	79
No	No	No	No	Yes	4
No	No	No	Yes	No	1
No	No	No	Yes	Yes	1
No	No	Yes	No	No	2
No	No	Yes	No	Yes	0
No	No	Yes	Yes	No	0
No	No	Yes	Yes	Yes	0
No	Yes	No	No	No	2
No	Yes	No	No	Yes	0
No	Yes	No	Yes	No	0
No	Yes	No	Yes	Yes	0
No	Yes	Yes	No	No	0
No	Yes	Yes	No	Yes	0
No	Yes	Yes	Yes	No	0
No	Yes	Yes	Yes	Yes	0
Yes	No	No	No	No	5
Yes	No	No	No	Yes	0
Yes	No	No	Yes	No	1
Yes	No	No	Yes	Yes	0
Yes	No	Yes	No	No	1
Yes	No	Yes	No	Yes	0
Yes	No	Yes	Yes	No	0
Yes	No	Yes	Yes	Yes	0
Yes	Yes	No	No	No	1
Yes	Yes	No	No	Yes	0
Yes	Yes	No	Yes	No	0
Yes	Yes	No	Yes	Yes	0
Yes	Yes	Yes	No	No	0
Yes	Yes	Yes	No	Yes	0
Yes	Yes	Yes	Yes	No	0
Yes	Yes	Yes	Yes	Yes	1

Notes:

To eliminate permutations that include missing data, this aspect of the analysis is based on the 752 subjects who provided information on oral disadvantage at baseline and at every follow-up during the 24-month observation period.

Pattern of self-reported oral disadvantage due to function.

Baseline	6-month	12-month	18-month	24-month	Percent of Sample with Each Pattern (n=757)
No	No	No	No	No	46
No	No	No	No	Yes	3
No	No	No	Yes	No	2
No	No	No	Yes	Yes	1
No	No	Yes	No	No	3
No	No	Yes	No	Yes	1
No	No	Yes	Yes	No	1
No	No	Yes	Yes	Yes	1
No	Yes	No	No	No	4
No	Yes	No	No	Yes	0
No	Yes	No	Yes	No	1
No	Yes	No	Yes	Yes	0
No	Yes	Yes	No	No	1
No	Yes	Yes	No	Yes	1
No	Yes	Yes	Yes	No	1
No	Yes	Yes	Yes	Yes	1
Yes	No	No	No	No	5
Yes	No	No	No	Yes	2
Yes	No	No	Yes	No	1
Yes	No	No	Yes	Yes	1
Yes	No	Yes	No	No	1
Yes	No	Yes	No	Yes	1
Yes	No	Yes	Yes	No	0
Yes	No	Yes	Yes	Yes	3
Yes	Yes	No	No	No	2
Yes	Yes	No	No	Yes	1
Yes	Yes	No	Yes	No	1
Yes	Yes	No	Yes	Yes	2
Yes	Yes	Yes	No	No	2
Yes	Yes	Yes	No	Yes	2
Yes	Yes	Yes	Yes	No	1
Yes	Yes	Yes	Yes	Yes	9

Notes:

To eliminate permutations that include missing data, this aspect of the analysis is based on the 757 subjects who provided information on oral disadvantage at baseline and at every follow-up during the 24-month observation period.